




BMJ Open Changing mortality trends in countries and cities of the UK: a population-based trend analysis

David Walsh ¹, Gerry McCartney ², Jon Minton,² Jane Parkinson ², Deborah Shipton,² Bruce Whyte¹

To cite: Walsh D, McCartney G, Minton J, *et al*. Changing mortality trends in countries and cities of the UK: a population-based trend analysis. *BMJ Open* 2020;**10**:e038135. doi:10.1136/bmjopen-2020-038135

► Prepublication history and additional material for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-038135>).

Received 28 February 2020
Revised 10 October 2020
Accepted 16 October 2020

ABSTRACT

Objectives Previously improving life expectancy and all-cause mortality in the UK has stalled since the early 2010s. National analyses have demonstrated changes in mortality rates for most age groups and causes of death, and with deprived populations most affected. The aims here were to establish whether similar changes have occurred across different parts of the UK (countries, cities), and to examine cause-specific trends in more detail.

Design Population-based trend analysis.

Participants/setting Whole populations of countries and selected cities of the UK.

Primary and secondary outcome measures European age-standardised mortality rates (calculated by cause of death, country, city, year (1981–2017), age group, sex and—for all countries and Scottish cities—deprivation quintiles); changes in rates between 5-year periods; summary measures of both relative (relative index of inequality) and absolute (slope index of inequality) inequalities.

Results Changes in mortality from around 2011/2013 were observed throughout the UK for all adult age groups. For example, all-age female rates decreased by approximately 4%–6% during the 1980s and 1990s, approximately 7%–9% during the 2000s, but by <1% between 2011/2013 and 2015/2017. Equivalent figures for men were 4%–7%, 8%–12% and 1%–3%, respectively. This later period saw increased mortality among the most deprived populations, something observed in all countries and cities analysed, and for most causes of death: absolute and relative inequalities therefore increased. Although similar trends were seen across all parts of the UK, particular issues apply in Scotland, for example, higher and increasing drug-related mortality (with the highest rates observed in Dundee and Glasgow).

Conclusions The study presents further evidence of changing mortality in the UK. The timing, geography and socioeconomic gradients associated with the changes appear to support suggestions that they may result, at least in part, from UK Government ‘austerity’ measures which have disproportionately affected the poorest.

INTRODUCTION

The recent (pre-COVID-19 pandemic) slowdown in improvement in life expectancy and mortality rates in the UK has been highlighted by researchers^{1 2} and media^{3 4} alike. Similar

Strengths and limitations of this study

- We examine recent changes in mortality in the UK in the context of much longer term trends: almost 40 years in most cases.
- Given the importance of urban health to national outcomes, we include the largest cities in Scotland, England (London excepted) and Northern Ireland.
- We analyse data for 10 major causes of death (not just all causes combined), and by country-specific and (for Scotland) city-specific deprivation levels.
- Limitations include the fact that interpretation of trends at city level can be problematic, given the fluctuation in rates.

changing trends have been observed in many other high-income countries, although the slowdown has been particularly marked in the UK and the USA.⁵ Other high-income countries with higher life expectancy have seen continued improvements.^{2 5}

Data from Scotland and England (92% of the UK population) have shown that these changing mortality patterns have been observed for almost all age groups and for most causes of death.^{6 7} Worryingly, increasing mortality rates among the most socioeconomically deprived populations have also been observed; as a result, inequalities in all-cause mortality have widened considerably since around 2012.^{7–9} An emerging body of UK^{1 10–13} and international^{14 15} work suggests the recent stalling is likely to be associated with the implementation from 2010 of UK Government ‘austerity’ measures—cuts to public services and social security—which have particularly affected the most vulnerable populations.

The principal aim of this project was to establish whether similar changing mortality trends (in terms of rates of improvement, causes of death and socioeconomic inequalities) have occurred ubiquitously across the UK. This included examining cause-specific



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

¹Glasgow Centre for Population Health, Glasgow, Scotland, UK

²Public Health Scotland, Glasgow, Scotland, UK

Correspondence to

Dr David Walsh;
david.walsh.2@glasgow.ac.uk

trends in more detail, and focusing on selected individual cities, given the importance of urban health to national outcomes.¹⁶

METHODS

Mortality and population data

Numbers of deaths by year of registration, age, sex, underlying cause, city and country for Scotland, England & Wales and Northern Ireland were obtained from, respectively, the National Records of Scotland (NRS), the Office for National Statistics and the Northern Ireland Statistics and Research Agency. Data were available for the following years: 1974–2017 (Scotland); 1981–2017 (England & Wales); and 1997–2017 (Northern Ireland).

Data were obtained for all-cause deaths, and for the following 10 major individual causes: respiratory disease; ischaemic heart disease (IHD); cerebrovascular disease; all malignant neoplasms; lung cancer (malignant neoplasm of trachea, bronchus and lung); intentional self-harm (including events of undetermined intent); external causes; motor vehicle traffic accidents (MVTAs); alcohol-related causes; and drug-related poisonings. Causes were defined by groups of International Classification of Diseases 8th Revision (ICD-8), ICD-9 and ICD-10 codes: these are listed in the online supplemental table 1. As stated in online supplemental table 1, the definition of external causes overlaps with other causes of death, that is, MVTAs, intentional self-harm, drug-related poisonings. ICD-9 codes were used for the years 1979–1999 in Scotland, and for 1981–2000 in England & Wales and Northern Ireland; ICD-10 codes were used for all later years. ICD-8 codes were used for 1974–1978 (Scotland only) but the data for those years are not presented here.

Matching population data by year, 5-year age group, sex, city and country were obtained from the same national statistical agencies.

Geography

Scotland, England & Wales (combined) and Northern Ireland were the countries used in the main analyses. For analysis by deprivation quintiles (discussed further below), England alone, rather than England & Wales, was used. With the exception of London, the largest cities in each country were selected: Glasgow, Edinburgh, Dundee and Aberdeen in Scotland; Liverpool, Manchester, Birmingham, Leeds, Sheffield and Bristol in England; and Belfast in Northern Ireland. London was excluded as its size and ethnic diversity makes meaningful comparisons with other cities problematic.¹⁷ Scottish and English cities were defined by current local authority boundaries. Belfast was defined by its 1992 local government district boundary.

Deprivation analyses employed the separate Scottish, English and Northern Irish area-based indices of deprivation: the Scottish Index of Multiple Deprivation (SIMD),¹⁸ the (English) Index of Multiple Deprivation (IMD)¹⁹ and the Northern Ireland Multiple Deprivation

Measure (NIMDM),²⁰ respectively. In all three cases data were available for the period 2001–2017. The SIMD has been updated multiple times: thus, the 2004 version was used for analyses covering the years 2001–2004, SIMD 2006 was used for 2005–2007, SIMD 2009 for 2008–2010, SIMD 2012 for 2011–2013 and SIMD 2016 for 2014–2017. Similarly, the (English) IMD 2004 was used for the years 2001–2005, IMD 2007 for 2006–2008, IMD 2010 for 2009–2013, IMD 2015 for 2014–2016 and IMD 2019 for 2017. For Northern Ireland, NIMDM 2010 was used for all years of analyses. Although there are differences in the spatial scale and the individual variables used in the construction of each nation's deprivation measure, all three share notable similarities in terms of their basic composition. The principal 'data domains' of each are effectively the same: income; employment; health; education, skills and training; crime; access to services; housing. In the Scottish index, housing is a separate category; in the English and Northern Irish indices it is instead contained within a 'living environment' domain. For all three measures of deprivation, similar methodologies are employed to calculate an overall index of relative deprivation, based on geographical area rankings across all data domains. Although the absolute values of the different indices cannot be directly compared, the similarity of composition and methodology associated with each provides helpful, and broadly comparable, overviews of inequality within each setting.

Statistical analyses

European age-standardised mortality rates per 100 000 population were calculated using the 2013 European Standard Population.²¹ Analyses were undertaken by sex, age (all ages, 0–64 years (the latter to examine premature deaths) and four broad groups across the life course: 0–14, 15–44, 45–64 and 65+ years), year, cause of death, city, country and deprivation quintile (see below). Three-year rolling average rates were derived; to quantify the rate of improvement over time, the percentage changes in rates between 3-year averages at 5-year intervals (ie, between 1981/1983 and 1985/1987, 1986/1988 and 1990/1992... up to 2011/2013 and 2015/2017) were calculated. Three-year averages were used to overcome the issue of fluctuating rates (especially at city level). For simplicity, we use the expression 'five year' interval to reflect the midpoints of the 3-year average (eg, 1982–1986 in relation to 1981/1983 to 1985/1987).

For the deprivation analyses, mortality rates by quintile were calculated on the basis of both national quintiles (based on levels of deprivation within individual countries) and—for Scottish cities only—city-specific quintiles (based on levels of deprivation within each individual city). City-specific quintiles are made publicly available by Information Services Division of NHS Scotland (now Public Health Scotland) for all versions of the SIMD used in the analyses, and were downloaded from their website. Equivalent data were not available for the English cities. In all analyses, quintile 1 represented the most deprived

fifth of the country's/city's small areas, and quintile 5 the least deprived fifth. The aim was to compare quintile rates within the same location (country or city), not between different locations. Note that as the English IMD does not include Wales, for comparison all-cause mortality rates by year and sex for England alone (rather than England & Wales combined) were also calculated. Analyses by deprivation were undertaken for all-cause deaths for all geographies, and for cause-specific mortality for all Scottish areas.

To examine changes in both absolute and relative inequalities in mortality (for the countries, Scottish cities and the causes of death), the slope index of inequality (SII) and the relative index of inequality (RII) respectively were calculated, based on the above deprivation quintiles. The SII calculates the gap in mortality rates across groups (here, the five quintiles), taking into account each quintile's rate (not just those of the least and most deprived), and each quintile's population size. The RII expresses this as a relative measure by dividing the SII by the rate for the whole population.²²

Analyses were undertaken using IBM SPSS Statistics V.25.

Patient and public involvement

Patients were not involved in this study.

RESULTS

Figure 1 presents all-cause standardised mortality rates for all ages by country and city for (A) males and (B) females between 1981/1983 and 2015/2017. At the national level, a change in the male death trend appears visible from around 2011/2013 for both Scotland and England, with

no or little improvement observed in the periods after that; there is greater fluctuation in rates in Northern Ireland. A change in rates appears more apparent, occurring slightly earlier, for deaths among females including those in Northern Ireland. At the city level, greater fluctuation in rates is clearly evident, as would be expected given the smaller population sizes and associated numbers of deaths: nonetheless, the majority of cities appear to have experienced a flattening, or worsening, of mortality rates in the last three to four time points shown.

To quantify the changes shown above, figure 2 shows the percentage change in rates between 5-year periods for Scotland, England & Wales and Northern Ireland. For women, mortality rates were decreasing by approximately 4%–6% during the 1980s and 1990s, with a faster improvement of approximately 7%–9% during the 2000s, and a much slower decrease of <1% between 2011/2013 and 2015/2017. Among men, mortality rates were decreasing slightly faster during the 1980s and early 1990s (by 4%–7%), with a faster improvement of approximately 8%–12% during the late 1990s and 2000s, and again a much slower decrease between 2011/2013 and 2015/2017 of 1%–3%. Similar data for the cities are shown in the online supplemental figure 1: these show a very similar overall pattern to the country-level analyses, although with greater fluctuation in the percentage change figures in some cities such as Dundee. However, it is also notable that—despite that fluctuation—in the most recent 5-year period mortality rates actually worsened among both males and females in Dundee and Aberdeen, and among females in Manchester, while there was virtually no improvement among males in Liverpool and females in Birmingham (–0.3% for both).




Figure 1 European age-standardised mortality rates (all ages, all causes) per 100 000 population, 3-year rolling averages, by UK country and city, 1981–2017, for (a) males and (b) females. EASR, European age-standardised mortality rate.




Figure 2 All-cause deaths, all ages: percentage change in standardised mortality rates between 5-year periods for (a) males and (b) females. n/a, not applicable.

Note that analyses based on more 'standard' 5-year periods (1981–1985, 1986–1990 up to 2011–2015) produced very similar results (data not shown).

Analyses of trends and changes in rates for ages 0–64 years (rather than all ages) are presented in online supplemental figure 2 for the countries of interest. While the results are broadly similar to those seen for all ages, the data suggest there has been a slight increase (rather than slowdown) in mortality rates for Scotland in the most recent period. Across the cities, a more notable change was observed in Dundee for males in this age group: a 26% increase in mortality between 2011/2013 and 2015/2017 (online supplemental figure 3). Notwithstanding the greater fluctuation in changes in rates in this age group at city level, it is also of note that there was an increase in female mortality rates in the last period in each English city.

Trends in all-cause rates for 45–64 years were similar to those described above for 0–64 years, and rates for 65+ years were broadly similar to those observed for all

ages. For the 15–44 age group, rates were notably higher in Scotland than England & Wales, and the mortality gap had become wider in the most recent period. These data are all shown in online supplemental figure 4(i). Although the principal focus of these analyses is adult mortality, online supplemental figure 4(ii) shows country-level rates over time for ages 0–14 years. It is notable that child mortality rates in Scotland have been consistently lower than in England & Wales since the late 1990s for females and early to mid-2000s for males, a reversal of the position at the beginning of the period. Rates are highest in Northern Ireland, although they are also subject to greater fluctuation over time.

Figure 3 again shows male all-cause mortality rates for all ages but additionally presents the rates for the least and most deprived deprivation quintiles within Scotland, England, Northern Ireland and an illustrative selection of three Scottish cities. Increased mortality rates are observed in the most recent period for the most deprived fifth of the population in each country/city




Figure 3 European age-standardised mortality rates (males, all ages, all causes) per 100 000 population, 3-year rolling averages by selected country, city and country/city-specific deprivation quintiles. EASR, European age-standardised mortality rate.

shown (although there is again more fluctuation in rates at the city level). Similar trends were observed for female mortality rates, as shown in online supplemental figure 5. Summary measures of inequalities (SIIs, RIIs) confirm widening absolute and relative inequalities across deprivation quintiles since 2011/2013 in all countries and cities analysed (online supplemental table 2a,b). For example, for all-cause deaths among men in England, the absolute gap across quintiles (as measured by the SII) increased from 738.8 in 2010/2012 to 784.6 in 2015/2017, despite having decreased between 2001/2003 and 2010/2012; relative inequalities increased over the whole period.

Mortality rates over time by deprivation quintile are explored in more detail for Scotland in figure 4, which presents data for females for all 10 causes of death analysed. A widening gap between the most and least

deprived quintiles in the most recent years of analysis can be seen for the majority of causes, in particular respiratory disease, external causes, drug-related poisonings and alcohol-related causes. Different trends are observed for suicide (where the gap has narrowed) and MVTAs (where numbers of deaths are relatively small and there is considerable fluctuation in rates across quintiles of deprivation). Generally, the same patterns, in terms of a widening gap between the most and least deprived deprivation quintiles for the majority of causes of death, are observed for male deaths (online supplemental figure 6). The main difference between the male and female trends relates to cancer mortality, in particular lung cancer, with a widening deprivation gap observed for females but not males. The general pattern of widening inequalities for the majority of causes is




Figure 4 European age-standardised mortality rates (females, all ages) per 100 000 population, 3-year rolling averages for 10 causes of death, Scotland and most and least deprived deprivation quintiles. EASR, European age-standardised mortality rate.




Figure 5 European age-standardised mortality rates (males, all ages) per 100 000 population, 3-year rolling averages, for all-cause deaths, ischaemic heart disease, alcohol-related causes and drug-related poisonings, UK countries and cities, 1981–2017. EASR, European age-standardised mortality rate.

confirmed by analysis of SIIs and RIIs (online supplemental table 3a,b).

Aside from evidence of a slowdown in mortality improvement and widening deprivation gaps (already shown in figures 1 and 3), the analyses of city-level mortality rates highlight a number of other issues. Foremost among them is that Glasgow stands out in terms of having the highest mortality rates of all the cities for all-cause deaths and the majority of causes analysed (including strikingly different trends for alcohol-related causes). Some exceptions do apply, however: for example, for all ages death rates from IHD are now marginally higher in Manchester for both males and females, while deaths from drug-related poisonings are now highest in Dundee. Some of these data are shown for males in figure 5, and for females in online supplemental figure 7.

Finally, figure 5/online supplemental figure 7 also highlight notable similarities and differences in long-term mortality trends between Scotland and England. For example: a narrowing of the gap between the countries for IHD; much higher drug-related poisonings in Scotland compared with England—although rates are increasing in both countries; and notably higher rates of death from alcohol-related causes in Scotland, with the most recent increase in rates much more pronounced in Scotland than

in England. The striking trends in drug-related poisonings are explored in more depth in online supplemental figure 8 (comparing the UK countries for selected age groups and by sex) and online supplemental figure 9 (showing data for Scotland and its most and least deprived quintiles for ages 0–64 years). There has been a widening divergence in death rates from this cause between Scotland and the rest of the UK over the period shown, and a further increase in rates is again observed since approximately 2012. Similar deprivation trends for drug-related poisonings are evident in the Scottish cities (data not shown).

DISCUSSION

Overall findings and implications

The study presents further evidence of a slowdown in mortality rate improvement over time within the UK. In some cases—deaths under 65 years in Scotland and all-age mortality in particular cities—rates have increased, rather than stalled, in recent years. These overall changes appear to be particularly driven by worsening mortality among the most socioeconomically disadvantaged populations. Similar trends are observed across all countries and cities in the UK; however, particular issues apply in Scotland, for example, in relation to drug-related

mortality. The timing, geographical coverage and socio-economic gradients associated with the changes appear to further support suggestions that recent changes in mortality are at least partly a consequence of UK Government 'austerity' measures.

Strengths and weaknesses

The analyses were based on data covering the whole population, not samples. Mortality is a robust population health indicator and is not subject to the limitations and potential biases associated with self-reported measures.^{23 24} We included the largest cities within Scotland and Northern Ireland and, with the exception of London, the six largest cities in England. We were able to examine recent changes in mortality in the context of much longer term trends: almost 40 years in the case of Scotland and England & Wales. We analysed data for a broad set of causes of death, not just all deaths combined.

The exclusion of London is arguably a weakness, although it was done to facilitate more meaningful comparisons across the other cities with more similar population sizes. Other limitations include the fact that time trend data for Northern Ireland were much more limited than for the rest of the UK. Interpretation of trends at city, rather than country level can also be problematic, given the fluctuation in rates that occur. Although the measures of area-based multiple deprivation that are employed within Scotland, England and Northern Ireland have many similarities, they are derived from different data sets and calculated at different spatial scales and are thus not directly comparable. We standardised mortality rates using 18 age groups (0–4 years to 85 years and above) while it is now recommended to standardise on the basis of 19 groups (0–4 years to 90 years and above)²⁵; however, the impact of this is fairly minimal. Although our youngest age bracket is 0–14 years, we did not include infant mortality (deaths under 1 year of age) as a distinct category: as a major focus of the work was city-based analyses, data on infant deaths were not requested from the various statistical agencies as numbers of deaths at that geographical level in the UK are very small; however, given recent evidence of increasing infant mortality rates in England (linked to increased child poverty rates), this was arguably an oversight.²⁶ Finally, the definition of drug-related poisonings is a broader, less sophisticated definition than that employed in official UK publications of drug-related mortality in the UK. In 2017, for example, there were 1037 such drug-related poisonings in Scotland, 11% higher than the 934 drug-related deaths recorded by the NRS.²⁷

Relevance to other studies

Slower improvement in mortality and life expectancy in the UK in recent years has been shown by various authors and organisations.^{1 2 5–8} Similar trends have been observed in a number of countries, although in a recent analysis of 20 high-income countries, only the USA had experienced a greater reduction in improvement than

the UK.⁵ As others have proposed, it appears increasingly likely that these changes are at least in part attributable to UK Government policy which since 2010 have brought about dramatic cuts to social security budgets and other public services, particularly affecting the most vulnerable.^{10–13} There is international evidence of the associations between such government measures and increased mortality rates,^{14 15} and the particular model of austerity adopted in the UK (based on spending cuts rather than a taxation approach) is known to be more regressive.¹³ UK research has highlighted associations between different UK austerity measures and increased child poverty,^{26 28} expansion of food banks,²⁹ increased homelessness,³⁰ poorer mental health among affected populations³¹ and, ultimately, increased numbers of deaths among the poorest at different ages.⁸ Interactions between such policy-driven factors and other influences such as high winter mortality in particular years have also been suggested.⁶

The widening socioeconomic inequalities in mortality since around 2012 have been shown previously for Scotland^{8 9} and England,^{7 32} but only at national level, and only for all-cause mortality and life expectancy. The increase in drug-related deaths within Scotland has been the focus of much media attention, resulting in two recent Westminster Parliamentary Committee enquiries,^{33 34} and the establishment of 'drugs death task force' by the Scottish Government in 2019.³⁵ The increase is known to be the result of a 'perfect storm' of factors: a previously described vulnerable cohort of drug users who are now encountering multiple morbidities as they age; increased affordability and accessibility of chosen drugs; and the aforementioned UK Government austerity measures which have impacted on both individual income and funding of drug-related and other relevant social services.^{33 34 36} These drug mortality trends are likely to have influenced the overall increasing death rates in Scotland (and in Dundee) among 0–64-year-olds.

Alcohol mortality trends have historically been impacted by changes in price and availability in combination with socioeconomic vulnerability,^{37 38} while the reasons for Glasgow's particularly high mortality rates (including from alcohol, drugs and other causes) have been described in detail previously, being attributable to higher socioeconomic deprivation alongside an additional vulnerability created by a combination of multiple historical factors including worse living conditions and adverse policymaking at different levels of government.³⁹

The contrasting trends in lung cancer mortality between males (decreasing rates) and females (increasing rates) living in the most deprived quintile in Scotland are also best explained in terms of different cohorts, with females having started smoking later than males: this has been shown in a number of other analyses.⁴⁰ In addition, one of the other notable differences between male and female trends—the apparent earlier slowdown in mortality improvement for females—is worthy of further analysis. Data presented here for both countries and cities suggest a

change in female mortality rates from around 2010/2012 or 2011/2013, potentially suggesting women's circumstances might be more sensitive to government austerity policies. However, more detailed analysis of Scottish data by Fenton *et al*² instead suggested a change between 2013 and 2014. Finally, although the main focus of the paper is adult mortality, the extent to which Scotland's child mortality rate has improved relative to that of England & Wales may be associated with a similar widening gap in child poverty levels between the countries over a similar time period is worthy of further investigation.⁴¹

CONCLUSIONS

These results add to the growing body of evidence of changing mortality rates within the UK in recent years and their likely political causes. With mortality rates rising among the UK's most deprived populations even prior to the COVID-19 pandemic, it is imperative that a range of policies are introduced to protect the health of the most vulnerable in society.

Twitter Gerry McCartney @gerrymccartney1 and Jon Minton @JonMinton

Acknowledgements We are grateful to the various individuals and organisations who supplied the required data: National Records of Scotland (NRS), the Office for National Statistics (ONS) and the Northern Ireland Statistics and Research Agency (NISRA). Particular thanks are due to Elaine Longden at NISRA for helpful advice and interpretation of trends, and also to Rebecca Holleyat at ONS. In addition, we gratefully acknowledge the work of the Geography, Population and Deprivation (GPD) Team of ISD Scotland for the provision of various SIMD deprivation lookup files.

Contributors DW originally conceived the study. The research questions and analysis plan were agreed by all authors (GMC, JM, JP, DS and BW). DW undertook the analyses and drafted the manuscript. All authors provided substantial critical input to improve the manuscript and all authors approved the final draft.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. All data relevant to the study are included in the article or uploaded as supplementary information. No additional data are available. The analysed data presented in the paper are not publicly available. However, the data on which the analyses were based can be requested from the national statistical agencies listed in the manuscript.

This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs


David Walsh <http://orcid.org/0000-0002-3390-5039>
Gerry McCartney <http://orcid.org/0000-0001-6341-3521>
Jane Parkinson <http://orcid.org/0000-0002-5697-1645>

REFERENCES


- Hiam L, Harrison D, McKee M, *et al*. Why is life expectancy in England and Wales 'stalling'? *J Epidemiol Community Health* 2018;72:404–8.
- Fenton L, Minton J, Ramsay J, *et al*. Recent adverse mortality trends in Scotland: comparison with other high-income countries. *BMJ Open* 2019;9:e029936.
- BBC. Life expectancy progress in UK 'stops for first time', 2018. Available: <https://www.bbc.co.uk/news/health-45638646>
- BBC. Scots' life expectancy falls for first time in 35 years, 2018. Available: <https://www.bbc.co.uk/news/uk-scotland-45636457>
- Ho JY, Hendi AS. Recent trends in life expectancy across high income countries: retrospective observational study. *BMJ* 2018;362:k2562.
- Ramsay J, Minton J, Fischbacher C, *et al*. How have changes in death by cause and age group contributed to the recent stalling of life expectancy gains in Scotland? Comparative decomposition analysis of mortality data, 2000–2002 to 2015–2017. *BMJ Open* 2020;10:e036529.
- Public Health England. A review of recent trends in mortality in England. London: Public Health England, 2018. Available: <https://www.gov.uk/government/publications/recent-trends-in-mortality-in-england-review-and-data-packs>
- Fenton L, Wyper GM, McCartney G, *et al*. Socioeconomic inequality in recent adverse all-cause mortality trends in Scotland. *J Epidemiol Community Health* 2019;73:971–4.
- Scottish Government. Long-term monitoring of inequalities. December 2018 report. Edinburgh: Scottish government, 2018. Available: <https://www.gov.scot/publications/long-term-monitoring-health-inequalities-december-2018-report>
- Dorling D. Austerity and mortality. In: Cooper V, Whyte D, eds. *The violence of austerity*. London: Pluto Press, 2017.
- Green M, Dorling D, Minton J. The geography of a rapid rise in elderly mortality in England and Wales, 2014–15. *Health Place* 2017;44:77–85.
- Loopstra R, McKee M, Katikireddi SV, *et al*. Austerity and old-age mortality in England: a longitudinal cross-local area analysis, 2007–2013. *J R Soc Med* 2016;109:109–16.
- Stuckler D, Reeves A, Loopstra R, *et al*. Austerity and health: the impact in the UK and Europe. *Eur J Public Health* 2017;27:18–21.
- Rajmil L, Fernández de Sanmamed M-J. Austerity policies and mortality rates in European countries, 2011–2015. *Am J Public Health* 2019;109:768–70.
- Toffolutti V, Suhrcke M. Does austerity really kill? *Econ Hum Biol* 2019;33:211–22.
- Galea S, Vlahov D. Urban health: evidence, challenges, and directions. *Annu Rev Public Health* 2005;26:341–65.
- Minton JW, McCartney G. Is there a north-south mortality divide in England or is London the outlier? *Lancet Public Health* 2018;3:e556–7.
- Scottish Government. Scottish index of multiple deprivation website. Available: <https://www2.gov.scot/Topics/Statistics/SIMD>
- Noble S, McLennan D, Noble M, *et al*. The English indices of deprivation 2019: research report. London: Ministry of housing, communities and local government, 2019. Available: <https://www.gov.uk/government/publications/english-indices-of-deprivation-2019-research-report>
- Northern Ireland Statistics and Research Agency (NISRA). Northern Ireland multiple deprivation measure 2010. Belfast: NISRA, 2010. Available: https://www.nisra.gov.uk/sites/nisra.gov.uk/files/publications/NIMDM_2010_Report_0.pdf
- European Commission. Revision of the European standard population - report of eurostat's task force. Luxembourg: publications office of the European union, 2013. Available: <https://ec.europa.eu/eurostat/documents/3859598/5926869/KS-RA-13-028-EN.PDF/e713fa79-1add-44e8-b23d-5e8fa09b3f8f>
- Regidor E. Measures of health inequalities: part 2. *J Epidemiol Community Health* 2004;58:900–3.
- Layes A, Asada Y, Keparat G. Whiners and deniers - what does self-rated health measure? *Soc Sci Med* 2012;75:1–9.
- Sen A. Health - perception versus observation: self-reported morbidity has severe limitations and can be extremely misleading. *BMJ* 2002;324:860–1.

- 25 Evans J. Implementing the 2013 European standard population: the impact of selected upper age limits on mortality statistics. London: office for national statistics, 2014. Available: <https://webarchive.nationalarchives.gov.uk/20160106020035/http://www.ons.gov.uk/ons/guide-method/user-guidance/health-and-life-events/revised-european-standard-population-2013--2013-esp-/impact-of-varying-the-2013-esp-upper-age-limit-on-mortality-statistics.doc>
- 26 Taylor-Robinson D, Lai ETC, Wickham S, *et al*. Assessing the impact of rising child poverty on the unprecedented rise in infant mortality in England, 2000-2017: time trend analysis. *BMJ Open* 2019;9:e029424.
- 27 National Records of Scotland (NRS). Drug-related deaths in Scotland in 2017. Edinburgh: NRS, 2018. Available: <https://www.nrscotland.gov.uk/files//statistics/drug-related-deaths/17/drug-related-deaths-17-pub.pdf>
- 28 Reed H, Stark G. *Forecasting child poverty in Scotland*. Edinburgh: Scottish Government, 2018.
- 29 Prayogo E, Chater A, Chapman S, *et al*. Who uses foodbanks and why? Exploring the impact of financial strain and adverse life events on food insecurity. *J Public Health* 2018;40:676-83.
- 30 Loopstra R, Reeves A, Barr B, *et al*. The impact of economic downturns and budget cuts on homelessness claim rates across 323 local authorities in England, 2004-12. *J Public Health* 2016;38:417-25.
- 31 Katikireddi SV, Molaodi OR, Gibson M, *et al*. Effects of restrictions to income support on health of lone mothers in the UK: a natural experiment study. *Lancet Public Health* 2018;3:e333-40.
- 32 Bennett JE, Pearson-Stuttard J, Kontis V, *et al*. Contributions of diseases and injuries to widening life expectancy inequalities in England from 2001 to 2016: a population-based analysis of vital registration data. *Lancet Public Health* 2018;3:e586-97.
- 33 UK Parliament Scottish Affairs Committee. Problem drug use in Scotland. London: UK Parliament, 2019. Available: <https://publications.parliament.uk/pa/cm201919/cmselect/cm��otaf/44/4402.htm>
- 34 UK Parliament Health and Social Care Committee. Drugs policy. London: UK Parliament, 2019. Available: <https://publications.parliament.uk/pa/cm201919/cmselect/cmhealth/143/14302.htm>
- 35 Scottish Government. Drug deaths task force, 2020. Available: <https://www.gov.scot/groups/drug-deaths-task-force>
- 36 Parkinson J, Minton J, Lewsey J, *et al*. Drug-related deaths in Scotland 1979-2013: evidence of a vulnerable cohort of young men living in deprived areas. *BMC Public Health* 2018;18:357.
- 37 McCartney G, Bouttell J, Craig N, *et al*. Explaining trends in alcohol-related harms in Scotland, 1991-2011 (I): the role of incomes, effects of socio-economic and political adversity and demographic change. *Public Health* 2016;132:13-23.
- 38 McCartney G, Bouttell J, Craig N, *et al*. Explaining trends in alcohol-related harms in Scotland 1991-2011 (II): policy, social norms, the alcohol market, clinical changes and a synthesis. *Public Health* 2016;132:24-32.
- 39 Walsh D, McCartney G, Collins C, *et al*. History, politics and vulnerability: explaining excess mortality in Scotland and Glasgow. *Public Health* 2017;151:1-12.
- 40 Wensink M, Alvarez J-A, Rizzi S, *et al*. Progression of the smoking epidemic in high-income regions and its effects on male-female survival differences: a cohort-by-age analysis of 17 countries. *BMC Public Health* 2020;20:39.
- 41 Joseph Rowntree Foundation (JRF). Uk poverty 2019/20. York: JRF, 2020. Available: <https://www.jrf.org.uk/report/uk-poverty-2019-20>


Web Figure 1. All cause deaths, all ages: percentage change in standardised mortality rates between five year periods – cities



Web Figure 2. European age-standardised mortality rates (0-64 years, all causes) per 100,000 population, three-year rolling averages, by UK country, 1981-2017; and percentage change in standardised mortality rates between five year periods




Web Figure 3. All cause deaths, 0-64 years: percentage change in standardised mortality rates between five year periods – cities




Web Figure 4. European age-standardised mortality rates (all causes) per 100,000 population, three-year rolling averages, by age group and UK country, 1981-2017 (note different axis scales on each chart)


Web Figure 4 (i) adult age groups




Web Figure 4 (ii) 0-14 years




Web Figure 5. European age-standardised mortality rates (females, all ages, all causes) per 100,000 population, three-year rolling averages by selected country, city and country/city-specific deprivation quintiles




Web Figure 6. European age-standardised mortality rates (males, all ages) per 100,000 population, three-year rolling averages for 10 causes of death, Scotland and most and least deprived deprivation quintiles




Web Figure 7. European age-standardised mortality rates (females, all ages) per 100,000 population, three-year rolling averages, for all-cause deaths, ischaemic heart disease, alcohol-related causes and drug-related poisonings, UK countries and cities, 1981-2017



Web Figure 8. European age-standardised mortality rates per 100,000 population, three-year rolling averages, for drug-related poisonings by selected age group, sex, and UK country 1981-2017



Web Figure 9. European age-standardised mortality rates per 100,000 population for drug-related poisonings, three-year rolling averages, Scotland and most and least by deprivation quintiles, 1981-2017



Web Table 1. Causes of death and associated ICD codes

Cause	ICD8 code(s)	ICD9 code(s)	ICD10 codes
Respiratory disease	460-519	460–519	J00-J99
Ischaemic heart disease	410-414	410-414	I20-I25
Cerebrovascular disease	430-438	430-438	I60-I69
All malignant neoplasms	140-209	140-208	C00-C97
Lung cancer (malignant neoplasm of trachea/bronchus/lung)	162	162	C33-C34
Intentional self-harm (including events of undetermined intent)	E950-E959, E980-E989	E950-E959, E980-E989	X60-X84, Y10-Y34, Y87.0, Y87.2
External causes	E800-E999	E800-E999	ICD10 V01-Y98
Motor vehicle traffic accidents (MVTAs) ⁱ	E810-E819	E810-E819	V02-V04, V09, V12-V14, V19-V79, V86-V89
Alcohol related causes ⁱⁱ	262,291,303,571.0, 571.9, E860	291, 303, 305.0, 425.5, 571.0-571.5, 571.8, 571.9, E860	F10, G312, G621, I426, K292, K70, K73, K740-K742, K746-K749, K860, X45, X65, Y15
Drug related poisonings ⁱⁱⁱ	304, E8530-E8539, E8540-E8549, E8550-E855.6, E8560-E8569, E950, E962, E980	304, 305.2-305.9, E850-E858, E950.0-E950.5, E9620, E980.0-E980.5	F11-F16, F18, F19, X40-X44, X60-X64, X85, Y10-Y14
Note that there are overlaps between a number of categories e.g.: all cancers and lung cancer; all external causes, intentional self-harm and drug-related poisonings; external causes and MVTAs			

ⁱ ICD9 and ICD10 codes as used in previous analyses e.g. Whyte B., Ajetunmobi T. Still 'The Sick Man of Europe'? Scottish mortality in a European context 1950-2010: an analysis of comparative mortality trends. Glasgow: GCPH; 2012

ⁱⁱ These are the groups of codes agreed by National Records of Scotland (NRS) and the (UK) Office for National Statistics (ONS) in 2007. They have since been updated, but the request for mortality data from national agencies pre-dated the change in codes.

ⁱⁱⁱ These are the set of codes previously deemed most comparable between Scotland and England & Wales. See: Walsh D., Bendel N., Jones R., Hanlon P. It's not 'just deprivation': Why do equally deprived UK cities experience different health outcomes? Public Health 2010; 124: 487-495.

Web Table 2a. Slope Index of Inequality (SII) and Relative Index of Inequality (RII) for all-cause mortality rates by deprivation quintile - males

	Scotland		England		Northern Ireland	
	SII	RII	SII	RII	SII	RII
2001-2003	1168.8	0.69	823.3	0.56	620.7	0.41
2002-2004	1166.5	0.70	821.8	0.58	536.0	0.36
2003-2005	1118.8	0.69	811.7	0.59	562.0	0.39
2004-2006	1102.6	0.71	796.5	0.60	593.3	0.42
2005-2007	1133.7	0.74	792.7	0.61	639.2	0.46
2006-2008	1129.6	0.76	797.3	0.63	653.7	0.47
2007-2009	1106.7	0.76	795.0	0.65	629.1	0.47
2008-2010	1044.7	0.74	776.0	0.65	615.4	0.47
2009-2011	1030.8	0.75	755.6	0.65	569.9	0.45
2010-2012	1018.7	0.76	738.8	0.65	586.9	0.48
2011-2013	1019.2	0.77	743.0	0.66	615.6	0.50
2012-2014	1012.4	0.78	745.6	0.67	609.3	0.51
2013-2015	1022.8	0.78	766.9	0.69	606.7	0.51
2014-2016	1050.7	0.81	778.2	0.71	616.5	0.53
2015-2017	1078.2	0.83	784.6	0.71	656.0	0.55

	Glasgow		Edinburgh		Dundee		Aberdeen	
	SII	RII	SII	RII	SII	RII	SII	RII
2001-2003	1373.7	0.64	1365.7	0.87	1023.4	0.57	942.5	0.58
2002-2004	1383.3	0.64	1292.4	0.84	993.9	0.56	983.6	0.62
2003-2005	1270.9	0.61	1198.3	0.79	877.8	0.51	999.4	0.66
2004-2006	1294.3	0.65	1229.5	0.84	778.1	0.49	1061.9	0.70
2005-2007	1279.0	0.65	1314.4	0.93	943.0	0.62	1076.5	0.71
2006-2008	1299.4	0.67	1265.2	0.93	1041.7	0.70	1207.9	0.82
2007-2009	1239.5	0.65	1193.2	0.90	1044.4	0.68	1200.6	0.83
2008-2010	1138.2	0.62	1068.5	0.81	912.2	0.61	1161.8	0.85
2009-2011	1187.1	0.67	1127.3	0.87	848.4	0.56	1061.8	0.80
2010-2012	1238.5	0.71	1045.7	0.82	986.1	0.66	905.0	0.70
2011-2013	1300.2	0.75	1081.5	0.87	1061.9	0.74	865.9	0.66
2012-2014	1266.5	0.75	1120.2	0.92	1235.2	0.88	844.7	0.64
2013-2015	1248.0	0.74	1187.1	0.97	1249.0	0.88	936.0	0.69
2014-2016	1278.6	0.76	1211.4	1.00	1422.1	0.97	1001.7	0.74
2015-2017	1329.8	0.78	1178.6	0.97	1443.5	0.96	1108.3	0.83

Web Table 2b. Slope Index of Inequality (SII) and Relative Index of Inequality (RII) for all-cause mortality rates by deprivation quintile - females

	Scotland		England		Northern Ireland	
	SII	RII	SII	RII	SII	RII
2001-2003	568.3	0.48	453.0	0.44	226.3	0.21
2002-2004	580.2	0.49	465.9	0.46	247.1	0.24
2003-2005	581.9	0.50	473.7	0.47	289.7	0.29
2004-2006	587.7	0.53	461.6	0.48	337.1	0.34
2005-2007	585.8	0.53	455.5	0.48	329.6	0.34
2006-2008	603.7	0.56	457.8	0.50	333.4	0.34
2007-2009	616.0	0.58	461.2	0.51	341.8	0.36
2008-2010	617.8	0.59	459.3	0.52	339.4	0.36
2009-2011	599.1	0.59	453.9	0.53	329.2	0.36
2010-2012	607.3	0.60	454.5	0.54	323.0	0.36
2011-2013	607.4	0.61	469.5	0.56	331.6	0.37
2012-2014	628.6	0.63	478.3	0.57	342.5	0.38
2013-2015	643.4	0.65	502.3	0.60	368.3	0.41
2014-2016	662.5	0.67	512.9	0.61	374.3	0.42
2015-2017	685.7	0.69	528.6	0.63	367.6	0.41

	Glasgow		Edinburgh		Dundee		Aberdeen	
	SII	RII	SII	RII	SII	RII	SII	RII
2001-2003	562.8	0.41	579.4	0.54	445.9	0.36	493.1	0.45
2002-2004	676.4	0.50	606.3	0.57	534.1	0.45	543.6	0.49
2003-2005	620.7	0.46	575.0	0.55	570.7	0.48	445.1	0.40
2004-2006	683.0	0.52	538.4	0.53	612.9	0.54	481.0	0.45
2005-2007	656.8	0.50	515.7	0.52	605.7	0.56	498.0	0.47
2006-2008	678.3	0.52	589.1	0.61	585.4	0.55	576.5	0.55
2007-2009	694.3	0.55	639.7	0.66	556.6	0.51	526.2	0.50
2008-2010	709.5	0.58	676.2	0.72	493.5	0.45	520.1	0.51
2009-2011	757.4	0.64	676.8	0.73	536.6	0.50	483.6	0.49
2010-2012	781.4	0.66	698.6	0.75	643.8	0.60	394.3	0.40
2011-2013	795.7	0.67	692.6	0.75	624.7	0.59	442.3	0.46
2012-2014	744.4	0.63	665.3	0.73	605.3	0.59	512.1	0.52
2013-2015	740.9	0.63	704.1	0.78	498.5	0.48	665.4	0.67
2014-2016	794.7	0.68	726.0	0.81	684.7	0.66	637.8	0.64
2015-2017	887.7	0.75	767.0	0.86	783.1	0.73	597.3	0.61

Web Table 3a. Slope Index of Inequality (SII) and Relative Index of Inequality (RII) for mortality rates by deprivation quintile: 10 major causes of death, Scotland, males

	Respiratory disease		Ischaemic heart disease		Cerebrovascular disease		All malignant neoplasms		Lung cancer	
	SII	RII	SII	RII	SII	RII	SII	RII	SII	RII
2001-2003	197.1	0.90	283.8	0.74	62.7	0.37	269.4	0.59	152.3	1.18
2002-2004	202.0	0.93	267.4	0.72	60.1	0.36	269.7	0.60	146.6	1.18
2003-2005	197.2	0.91	256.1	0.73	61.6	0.39	262.1	0.60	147.4	1.22
2004-2006	183.9	0.90	225.0	0.69	52.2	0.36	276.9	0.64	152.3	1.29
2005-2007	185.4	0.90	233.0	0.76	50.0	0.37	289.5	0.68	158.0	1.33
2006-2008	189.1	0.93	215.6	0.76	53.0	0.41	280.1	0.67	155.7	1.35
2007-2009	196.8	0.98	211.3	0.78	55.2	0.45	268.9	0.65	154.6	1.36
2008-2010	189.4	0.99	190.2	0.75	50.9	0.44	254.0	0.63	146.6	1.34
2009-2011	185.0	1.01	182.7	0.77	45.7	0.43	268.0	0.67	144.3	1.33
2010-2012	177.8	1.00	187.1	0.83	39.9	0.40	270.8	0.68	140.1	1.33
2011-2013	184.0	1.05	180.3	0.85	41.3	0.43	269.7	0.68	143.2	1.39
2012-2014	174.4	1.02	182.7	0.90	34.7	0.38	262.5	0.68	140.8	1.41
2013-2015	179.8	1.05	174.6	0.89	39.0	0.44	253.6	0.66	136.3	1.40
2014-2016	178.9	1.06	169.4	0.90	42.0	0.49	255.9	0.67	127.8	1.35
2015-2017	185.3	1.12	165.6	0.89	45.1	0.54	254.3	0.67	126.4	1.38

	External causes		Drug related poisonings		Alcohol-related causes		Suicide (incl. undetermined)		Motor vehicle traffic accidents	
	SII	RII	SII	RII	SII	RII	SII	RII	SII	RII
2001-2003	77.6	1.16	37.5	2.58	106.5	2.33	35.7	1.4	1.0	0.10
2002-2004	79.3	1.17	36.6	2.47	110.8	2.40	34.3	1.4	-0.6	-0.07
2003-2005	72.7	1.10	32.1	2.36	106.3	2.33	29.4	1.3	-1.0	-0.11
2004-2006	72.9	1.11	36.0	2.49	106.9	2.40	30.1	1.3	1.5	0.16
2005-2007	69.7	1.11	39.4	2.50	102.7	2.40	29.8	1.3	2.4	0.29
2006-2008	69.4	1.11	47.0	2.53	102.0	2.46	31.8	1.3	3.1	0.38
2007-2009	67.1	1.11	48.3	2.49	91.5	2.40	31.9	1.4	1.8	0.25
2008-2010	64.8	1.11	48.4	2.48	87.3	2.35	31.6	1.4	1.3	0.20
2009-2011	72.9	1.17	46.3	2.43	76.5	2.20	32.8	1.5	2.1	0.36
2010-2012	80.0	1.21	47.1	2.46	72.6	2.18	31.3	1.3	1.5	0.28
2011-2013	89.7	1.28	47.5	2.47	65.0	2.11	30.4	1.3	2.7	0.51
2012-2014	89.8	1.32	49.7	2.56	66.4	2.20	25.9	1.2	2.0	0.38
2013-2015	89.5	1.33	51.4	2.59	64.3	2.13	22.9	1.1	2.7	0.53
2014-2016	101.0	1.43	58.7	2.66	67.4	2.14	22.7	1.2	2.9	0.57
2015-2017	112.4	1.50	65.9	2.71	67.0	2.08	23.5	1.2	2.9	0.65

Web Table 3b. Slope Index of Inequality (SII) and Relative Index of Inequality (RII) for mortality rates by deprivation quintile: 10 major causes of death, Scotland, females

	Respiratory disease		Ischaemic heart disease		Cerebrovascular disease		All malignant neoplasms		Lung cancer	
	SII	RII	SII	RII	SII	RII	SII	RII	SII	RII
2001-2003	110.5	0.75	135.7	0.64	24.3	0.15	154.0	0.52	84.3	1.24
2002-2004	116.7	0.78	125.1	0.61	24.8	0.16	160.7	0.55	87.5	1.26
2003-2005	121.1	0.80	119.8	0.62	22.6	0.15	157.1	0.54	86.5	1.23
2004-2006	124.3	0.83	111.9	0.62	19.5	0.14	155.5	0.54	88.7	1.23
2005-2007	127.1	0.84	109.4	0.66	10.2	0.08	155.4	0.54	93.3	1.28
2006-2008	133.5	0.88	103.8	0.67	17.6	0.14	158.2	0.55	96.2	1.29
2007-2009	136.3	0.92	102.6	0.71	18.5	0.16	169.6	0.60	99.7	1.34
2008-2010	137.8	0.96	96.2	0.71	24.5	0.22	174.3	0.62	98.4	1.32
2009-2011	137.4	1.01	87.4	0.69	22.2	0.21	175.6	0.63	99.7	1.35
2010-2012	147.2	1.09	80.9	0.67	22.7	0.23	173.9	0.62	104.2	1.40
2011-2013	146.3	1.10	76.1	0.67	20.3	0.21	174.7	0.63	107.5	1.45
2012-2014	143.6	1.10	83.9	0.79	19.5	0.21	183.7	0.66	109.6	1.49
2013-2015	139.5	1.06	84.9	0.83	23.9	0.27	186.3	0.68	106.7	1.49
2014-2016	146.9	1.12	87.4	0.90	24.0	0.29	184.8	0.69	105.6	1.51
2015-2017	150.8	1.14	80.7	0.86	25.5	0.32	180.6	0.68	106.7	1.53

	External causes		Drug related poisonings		Alcohol-related causes		Suicide (incl. undetermined)		Motor vehicle traffic accidents	
	SII	RII	SII	RII	SII	RII	SII	RII	SII	RII
2001-2003	17.0	0.50	13.2	2.26	35.9	1.96	12.5	1.47	0.1	0.04
2002-2004	18.1	0.53	12.7	2.25	35.2	1.94	11.5	1.39	0.2	0.06
2003-2005	18.7	0.56	12.0	2.10	35.0	1.91	10.9	1.34	0.1	0.02
2004-2006	21.7	0.68	12.6	2.17	35.4	1.90	10.1	1.34	0.5	0.19
2005-2007	23.5	0.76	12.6	2.28	35.6	1.93	11.6	1.57	0.7	0.27
2006-2008	25.2	0.84	14.6	2.50	34.5	1.95	12.1	1.64	0.4	0.14
2007-2009	22.7	0.74	16.8	2.55	30.6	1.84	11.3	1.48	0.1	0.04
2008-2010	21.1	0.70	17.9	2.49	29.0	1.80	9.6	1.30	-0.4	-0.17
2009-2011	23.4	0.74	18.7	2.39	26.5	1.66	9.7	1.25	0.0	0.02
2010-2012	29.6	0.92	19.5	2.41	25.7	1.77	10.5	1.31	-0.5	-0.26
2011-2013	32.9	1.01	19.0	2.35	23.1	1.68	10.0	1.27	-0.8	-0.49
2012-2014	31.8	0.99	19.3	2.47	21.8	1.70	9.2	1.28	-0.8	-0.48
2013-2015	32.7	0.97	21.1	2.52	23.0	1.74	7.8	1.12	-0.2	-0.10
2014-2016	37.6	1.03	27.1	2.71	24.2	1.78	7.5	1.04	0.3	0.20
2015-2017	46.1	1.18	31.3	2.69	24.3	1.77	6.6	0.98	0.6	0.33